

A notion of graph likelihood and an infinite monkey theorem

Christopher Banerji,^{1,2} Toufik Mansour,³ and Simone Severini⁴

¹*Department of Computer Science, and Centre of Mathematics & Physics in the Life Sciences and Experimental Biology, University College London, London WC1E 6BT, United Kingdom*

²*Statistical Cancer Genomics, Paul O’Gorman Building, UCL Cancer Institute, University College London, London WC1E 6BT, United Kingdom*

³*Department of Mathematics, University of Haifa, Haifa 31905, Israel*

⁴*Department of Computer Science, and Department of Physics & Astronomy, University College London, WC1E 6BT London, United Kingdom*

We play with a graph-theoretic analogue of the folklore infinite monkey theorem. We define a notion of graph likelihood as the probability that a given graph is constructed by a monkey in a number of time steps equal to the number of vertices. We present an algorithm to compute this graph invariant and closed formulas for some infinite classes. We have to leave the computational complexity of the likelihood as an open problem.

I. INTRODUCTION

The *infinite monkey theorem* is part of the popular culture [3]. A monkey sits in front to a typewriter hitting random keys. The probability that the monkey will type any given text tends to one, as the amount of time the monkey spends on the typewriter tends to infinity. The usual example is of Shakespeare’s Hamlet. Of course, the term “monkey” can refer to some abstract device producing random strings of symbols (*e.g.*, zeros and ones).

In this note, we consider an infinite monkey theorem, but for graphs rather than strings. Our setting involves a device which performs “non-preferential” attachment [2]. At time step $t + 1$, a new vertex is added to a graph G_t – the process starts from the single vertex graph, G_1 . The degree and the neighbours of the newly added vertex at step $t + 1$ are both chosen at random. The degree of the vertex is then $k \in \{0, 1, \dots, t\}$ and its neighbours are k random vertices in G_t . The (in fact obvious) analogue of the infinite monkey theorem is that every graph can be constructed in this way: the probability that the monkey will construct a given graph tends to one, as the amount of time the monkey spends on the “graphwriter” tends to infinity. Notice that the monkey makes two random choices, but these can be seen as a single one. Also, notice that the theorem is indeed a corollary of the usual infinite monkey theorem since we could encode a graph in a string (for example, by vectorizing the adjacency matrix). Below is a monkey enjoying the construction of the Petersen graph:



The construction is basically an excuse to discuss a graph invariant which we call *(graph) likelihood*. This is the probability that a given graph on t vertices is obtained by the construction after exactly t steps. In other words,

this is the probability that a monkey constructs a given graph on t vertices in exactly t seconds, assuming that the monkey adds a new vertex each second. For a string, this would correspond to the probability that the monkey types a given text in a time equal to the length of the string produced.

The likelihood is a plausible measure to quantify how difficult is to construct a graph in the way we propose. Intuitively, graphs with more symmetries have generally smaller likelihood. We will show as expected that bounds on the likelihood can be given in terms of the automorphism group. Specifically, the likelihood can not be larger than the reciprocal of the number of automorphisms. Graphs with trivial automorphism group are then potentially the ones admitting highest likelihood. We will describe an algorithm to compute the likelihood of a given graph. The algorithm uses a rooted tree decomposition which takes into account all possible ways to construct the graph by adding one vertex at the time. The algorithm suggests a closed formula for the likelihood.

The reminder of the paper is organized as follows. In Section II, we define the likelihood. In Section III, we give closed formulas for complete graphs, star graphs, paths, and cycles. In Section IV, we describe the algorithm. Section V lists some open problems. In particular, we could not to prove the complexity of computing the likelihood. The paper is practically self-contained.

II. GRAPH LIKELIHOOD

As usual, $G = (V, E)$ denotes a *graph*: $V(G) = \{v_1, v_2, \dots, v_t\}$ is a set whose elements are called *vertices* and $E(G) \subseteq V(G) \times V(G) - \{\{v_i, v_i\} : v_i \in V(G)\}$ is a set whose elements are called *edges*. The graph with a single vertex and no edges is denoted by K_1 . Our main object of study will be the construction given in the following definition. This is a special case of a construction already presented in [2].

Definition 1 (Construction) *We construct a graph $G_t = (V, E)$, starting from $G_1 = K_1$. The construction involves an iteration with discrete steps. At the t -th step of the iteration, the graph G_{t-1} is transformed into the graph G_t . The t -th step of the iteration is divided into three substeps:*

1. *We select a number $k \in \{0, 1, \dots, t-1\}$ with equal probability.
Assume that we have selected k .*
2. *We select k vertices of G_{t-1} with equal probability.
Assume that we have selected the vertices $v_1, v_2, \dots, v_k \in V(G_{t-1})$.*
3. *We add a new vertex t to G_{t-1} and the edges $\{v_1, t\}, \{v_2, t\}, \dots, \{v_k, t\} \in E(G_t)$.*

On the basis of the construction, the following definition is natural:

Definition 2 (Graph likelihood) *Let G be a graph on t vertices. The (graph) likelihood of G , denoted by $\mathcal{L}(G)$, is defined as the probability that $G_t = G$, where G_t is the graph given by the construction in Definition 1:*

$$\mathcal{L}(G) := \Pr[G_t = G].$$

For clarifying this notion, in the next section, we write closed formulas for the likelihood of graphs in some infinite families. We use rather uninteresting proof techniques, but these serve the purpose, at least for very simple graphs.

III. EXAMPLES

The *complete graph* K_t is defined as the graph on t vertices and $t(t-1)/2$ edges.

Proposition 3 *Let K_t be the complete graph on t vertices. Then, $\mathcal{L}(K_t) = 1/t!$.*

Proof. For K_t , the only significant step of the construction is the first one (*i.e.*, the selection of a number $k \in \{0, 1, \dots, t-1\}$ with equal probability). Therefore, $\mathcal{L}(K_t) = \prod_{i=1}^t \frac{1}{i}$. This equals $1/t!$ by definition. ■

The *star graph* $K_{1,t-1}$ is defined as the graph on t vertices, $v_1, v_2, \dots, v_t$, and the edges $\{v_1, v_2\}, \{v_1, v_3\}, \dots, \{v_1, v_{t-1}\}$. In a graph $G = (V, E)$, the *degree* of a vertex $i \in V(G)$ is defined and denoted by $d(i) = |\{j : \{i, j\} \in E(G)\}|$.

Proposition 4 *Let $K_{1,t-1}$ be the star graph on t vertices. Then, $\mathcal{L}(K_{1,t-1}) = \frac{t}{(t!)^2} \sum_{i=0}^{t-1} i!$.*

Proof. The star graph $K_{1,t-1}$ has 1 vertex of degree $t-1$ and $t-1$ vertices of degree 1. There are three cases relevant to the construction of $K_{1,t-1}$ such that $G_t = K_{1,t-1}$:

1. Suppose we add $t-1$ vertices, $1, 2, \dots, t-1$, of degree 0. At time t , we add a vertex, t , of degree $t-1$. Since $\Pr[d(i) = 0] = 1/i$, for $i = 1, 2, \dots, t-1$, and $\Pr[d(t) = t-1] = 1/t$, $\Pr[G_t = K_{1,t-1} \text{ by (1)}] = \prod_{i=1}^{t-1} \Pr[d(i) = 0] \cdot \Pr[d(t) = t-1] = \left(\prod_{i=1}^{t-1} \frac{1}{i}\right) \cdot \frac{1}{t} = \frac{1}{t!}$.
2. Suppose there is an edge $\{1, 2\} \in G_2$. Since $\Pr[d(3) = 1] = 1/3$, we distinguish two cases:
 - (a) $\Pr[\{1, 3\} \in E(G_3)] = 1/2$: $\Pr[d(2) = 1] \cdot \Pr[d(3) = 1] \cdot \Pr[\{1, 3\} \in E(G_3)] = \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{2} = \frac{1}{12}$. If $\{1, 3\} \in E(G_3)$ then each other edge of G_t , with $t \geq 4$, must be of the form $\{1, 4\}, \dots, \{1, t\}$ and $\Pr[\{1, t\} \in E(G_3)] = \frac{1}{t} \cdot \frac{1}{t-1}$.
 - (b) $\Pr[\{2, 3\} \in E(G_3)] = 1/2$: $\Pr[d(2) = 1] \cdot \Pr[d(3) = 1] \cdot \Pr[\{2, 3\} \in E(G_3)] = \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{2} = \frac{1}{12}$. If $\{2, 3\} \in E(G_3)$ then the situation is analogous to the previous case.

By combining together (a) and (b), it follows that

$$\Pr[G_t = K_{1,t-1} \text{ by (2)}] = 2 \prod_{i=2}^t \left(\frac{1}{i} \cdot \frac{1}{i-1} \right) = \frac{2}{t!(t-1)!}.$$

3. Suppose we add $k-1$ vertices, $1, 2, \dots, k-1$, of degree 0, where $k \geq 3$. At time k , we add a vertex, k , of degree $k-1$. Since $\Pr[d(i) = 0] = 1/i$, for $i = 1, 2, \dots, k-1$, and $\Pr[d(k) = k-1] = 1/k$,

$$\Pr[G_k = K_{1,k-1} \text{ by (3)}] = \prod_{i=1}^{k-1} \Pr[d(i) = 0] \cdot \Pr[d(k) = k-1] = \left(\prod_{i=1}^{k-1} \frac{1}{i} \right) \cdot \frac{1}{k} = \frac{1}{k!}.$$

The remaining $t-k$ vertices, $k+1, k+2, \dots, t$, must be of the form $\{k, k+1\}, \{k, k+2\}, \dots, \{k, t\}$ and

$$\Pr[\{k, k+j\} \in E(G_{k+j})] = \frac{1}{k+j} \cdot \frac{1}{k+j-1},$$

for each $j = 1, 2, \dots, t-k$. Hence,

$$\Pr[G_t = K_{1,t-1} \text{ by (3)}] = \sum_{k=3}^{t-1} \frac{1}{k!} \prod_{i=k+1}^t \left(\frac{1}{i} \cdot \frac{1}{i-1} \right).$$

The analysis carried out with the three cases above is sufficient to obtain the following formula:

$$\begin{aligned} \mathcal{L}(K_{1,t-1}) &= \Pr[G_t = K_{1,t-1} \text{ by (1)}] + \Pr[G_t = K_{1,t-1} \text{ by (3)}] + \Pr[G_t = K_{1,t-1} \text{ by (2)}] \\ &= \frac{1}{t!} + \frac{2}{t!(t-1)!} + \sum_{k=3}^{t-1} \frac{1}{k!} \prod_{i=k+1}^t \frac{1}{i(i-1)} = \frac{t}{(t!)^2} \sum_{i=0}^{t-1} i!. \end{aligned}$$

■

IV. COMPUTATION OF THE LIKELIHOOD

Is the likelihood defined for any graph? The answer is “yes”, as demonstrated by the next statement. This is a plausible graph-theoretic analogue of the infinite monkey theorem:

Proposition 5 *Any graph can be obtained with the construction in Definition 1.*

Proof. An *orientation* of G is a function $\alpha : E(G) \rightarrow E^+(G)$, where $E^+(G)$ is a set whose elements, called *arcs*, are ordered pairs of vertices such that either $\alpha(\{i, j\}) = (i, j)$ or $\alpha(\{i, j\}) = (j, i)$, for each $\{i, j\} \in E(G)$. An orientation is *acyclic* if it does not contain any directed cycles, *i.e.*, distinct vertices $v_1, \dots, v_k$ such that $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k), (v_k, v_1)$ are arcs. Clearly, every graph has an acyclic orientation. Every acyclic orientation determines at least one linear ordering $v_1 < v_2 < \dots < v_n$ of the vertices such that, for each edge $\{v_i, v_j\}$, we have $\alpha(\{v_i, v_j\}) = (v_i, v_j)$ if and only if $v_i < v_j$. This is also called a *topological ordering* of the vertices relative to the orientation. For a graph G , let $V(G) = \{w_1, w_2, \dots, w_t\}$ and let $w_1 < w_2 < \dots < w_t$ realize a topological ordering. We can always obtain $G_t = G$, if in the iteration we have $v_1 = w_1, v_2 = w_2, \dots, v_t = w_t$. ■

And, of course:

Proposition 6 *Every graph on n vertices has a positive likelihood. (More formally, $\mathcal{L}(G) > 0$ for any graph G .)*

Proposition 5 suggests a natural computational problem:

Problem 7 (Likelihood computation) *Given:* A graph G . *Task:* Compute $\mathcal{L}(G)$.

There are surely many ways to approach this problem. We consider an algorithm based on a tree whose vertices represent all intermediate graphs obtained during the construction.

Definition 8 (Identity representation) *Let $G = (V, E)$ be a graph on the set of vertices $V(G) = \{v_1, v_2, \dots, v_t\}$. Let us fix an arbitrary labeling of the vertices of G by a bijection $f : V(G) \rightarrow \{1, 2, \dots, t\}$. Once fixed the bijection, let us label the first row (resp. column) of the adjacency matrix of G , $A(G)$, by the number t , the second one by $t-1, \dots$, the last one by 1. The bijection f can be then represented by the ordered set $(1, 2, \dots, t)$. We can then define an acyclic orientation of the edges such that $\alpha(\{i, j\}) = (i, j)$ if and only if $i < j$, with $i, j = 1, 2, \dots, t$. The topological ordering relative to the orientation defines $G_1 = (1, \emptyset)$, $G_2 = (\{1, 2\}, E(G_2))$, ..., $G_t = (\{1, 2, \dots, t\}, E(G_t)) = G$. The pair $(A(G), (1, 2, \dots, t))$ given by the adjacency matrix $A(G)$ together with the ordered set $\text{id} := (1, 2, \dots, t)$ is said to be the identity representation of G .*

Remark 9 *The identity representation is arbitrary, since it entirely depends on the bijection f .*

A permutation of length n is a bijection $p : \{1, 2, \dots, t\} \rightarrow \{1, 2, \dots, t\}$. Hence, each permutation p corresponds to an ordered set $(p(1), p(2), \dots, p(t))$. The set of all permutations of length t is denoted by S_t . A permutation matrix P induced by a permutation p of length t is an $t \times t$ matrix such that $[P]_{i,j} = 1$ if $p(i) = j$ and $[P]_{i,j} = 0$, otherwise. Lower case letters denote permutations; upper case letters their induced matrices.

Definition 10 ((Generic) Representation) *Let $G = (V, E)$ be a graph on t vertices. Let $(A(G), \text{id})$ be the identity representation of G . The pair $(PA(G)P^T, p)$, where P is a permutation matrix induced by the permutation p is said to be a representation of G . A representation $(PA(G)P^T, p)$ is also denoted by $A_p(G)$.*

An automorphism of a graph $G = (V, E)$ is a permutation $p : V(G) \rightarrow V(G)$ such that $\{v_i, v_j\} \in E(G)$ if and only if $\{p(v_i), p(v_j)\} \in E(G)$. The set of all automorphisms of G , with the operation of composition of permutations “ $\circ$ ”, is a permutation group denoted by $\text{Aut}(G)$. Such a group is the full automorphism group of G . The permutation matrices P , induced by the elements of $\text{Aut}(G)$, are precisely the matrices such that $PA(G)P^T = A(G)$, i.e., $PA(G) = A(G)P$.

Lemma 11 *Let $G = (V, E)$ be a graph on t vertices. The total number of different representations of G is $t! / |\text{Aut}(G)|$.*

Proof. Let $A_{\text{id}}(G)$ be an identity representation of G . By the definition of full automorphism group, for each permutation $p \in \text{Aut}(G)$, we have $A_p(G) = PA_{\text{id}}(G)P^T = A_{\text{id}}(G)$. Let $q \in S_t - \text{Aut}(G)$. Then, there is a unique permutation $r \in S_t - \text{Aut}(G)$ such that $q = p \circ r$. It follows that $QA_{\text{id}}(G)Q^T = PRA_{\text{id}}(G)R^T P^T = PA_r(G)P^T = A_r(G)$. This indicates that each representation of G belongs to an equivalence class of representations. Since $|S_t| = t!$, the total number of different representations of G , i.e., the total number of equivalence classes of representations, is $t! / |\text{Aut}(G)|$. ■

Remark 12 *In the language of elementary group theory, the equivalence classes are the (left) cosets of the subgroup $\text{Aut}(G)$ in S_t .*


In order to design an algorithm for $\mathcal{L}(G)$, we need some further definitions. A subgraph $H = (V', E')$ of a graph $G = (V, E)$ is a graph such that $V' \subseteq V$ and $E' \subseteq E$. We say that a graph G contains a graph H if there is a subgraph of G isomorphic to H .

Definition 13 *Let G be any nonempty graph with t vertices, a path construction of G is a sequence $(H_1, H_2, \dots, H_t)$ of t graphs such that H_i has i vertices, $i = 1, 2, \dots, t$, and $H_i \subset H_{i+1}$, for each $i = 1, 2, \dots, t-1$; moreover, $H_t \cong G$. We denote the set of all path constructions of a graph G by $\text{Path}(G)$.*

It is clear that each path construction corresponds to an equivalence class of representations.

The set of path constructions can be represented as a rooted tree T_G as follows:

- The root of T_G is T_1 . This is the empty graph with a single vertex.
- Assume we already have all the vertices at level i (the level of the root is taken to be 1) in the tree T_G . Let $(T_1, T_2, \dots, T_i)$ be a path in T_G , if there exists a path construction $L = (T_1, T_2, \dots, T_i, H_{i+1}, \dots, H_t) \in \text{Path}(G)$ then we define H_{i+1} to be one of the children of the node T_i in T_G .



Example 14 The rooted tree T_{P_3} is given by

The above figure shows that the set of path constructions of P_3 is given by

$$\text{Path}(P_3) = \{(T_1, T_{21}, T_{31}), (T_1, T_{22}, T_{32}), (T_1, T_{22}, T_{33})\}.$$

Let $P = (H_1, H_2, \dots, H_t) \in \text{Path}(G)$ be any path construction of G . Fix i , then H_{i+1} is obtained by adding a vertex v_{i+1} of degree $d_{i+1}(P)$ to the graph H_i . Hence

$$\Pr[G_t = G, P \text{ is a path construction of } G] = \prod_{i=1}^t \frac{1}{i \binom{i-1}{d_i(P)}} = \frac{1}{t! \prod_{i=1}^t \binom{i-1}{d_i(P)}}.$$

From this algorithm, we obtain a relation between $\mathcal{L}(C_n)$ and $\mathcal{L}(P_n)$ as follows. Recall that C_n is the cycle on n vertices and P_n is the path on n vertices.

Corollary 15 For all $n \geq 3$, $\mathcal{L}(C_n) = \mathcal{L}(P_{n-1})/n \binom{n-1}{2}$.

An algorithm for computing $\mathcal{L}(G)$ can be based on the following theorem:

Theorem 16 Let G be a graph on t vertices. Then

$$\mathcal{L}(G) = \sum_{P \in \text{Path}(G)} \frac{1}{t! \prod_{i=1}^t \binom{i-1}{d_i(P)}}.$$

A simple example is useful:

Example 17 Let P_3 be the path graph on 3 vertices. By Example 14, we find that

$$\begin{aligned} \Pr[G_t = G, (T_1, T_{21}, T_{31}) \text{ is a path construction of } G] &= 1 \cdot \frac{1}{2} \cdot \frac{1}{3} = \frac{1}{6}, \\ \Pr[G_t = G, (T_1, T_{22}, T_{32}) \text{ is a path construction of } G] &= 1 \cdot \frac{1}{2} \cdot \frac{1}{3 \cdot 2} = \frac{1}{12}, \\ \Pr[G_t = G, (T_1, T_{22}, T_{33}) \text{ is a path construction of } G] &= 1 \cdot \frac{1}{2} \cdot \frac{1}{3 \cdot 2} = \frac{1}{12}. \end{aligned}$$

Then, $\mathcal{L}(G) = \frac{1}{6} + \frac{1}{12} + \frac{1}{12} = \frac{1}{3}$.

By Theorem 16 and the fact that $|\text{Path}(G)|$ is exactly equal to the number of representations of G , i.e. $|\text{Path}(G)| = t! / |\text{Aut}(G)|$, we obtain the following bounds:

Corollary 18 Let G be any nonempty graph on t vertices. Then

$$\frac{1}{|\text{Aut}(G)| \prod_{i=1}^t \binom{i-1}{\lfloor (i-1)/2 \rfloor}} \leq \mathcal{L}(G) \leq \frac{1}{|\text{Aut}(G)|}.$$

We give two general examples:

Example 19 Let G be a graph on t vertices with exactly s edges incident with $2s$ vertices. Any path P of $\text{Path}(G)$ can be seen as a path from a single vertex to the graph G . At levels $i_1, i_2, \dots, i_s$, we have added an edge between the new vertex and a vertex of degree zero. In all other levels we just added a new vertex. Therefore,

$$\mathcal{L}(G) = \frac{1}{t!} \sum_{2 \leq i_1 < i_2 < \dots < i_s \leq t} \prod_{j=1}^s \frac{i_j + 1 - 2j}{i_j - 1}.$$

Example 20 Let G be a graph on t vertices with exactly one edge, then

$$\mathcal{L}(G) = \frac{1}{t!} \sum_{i=2}^t 1 = \frac{t-1}{t!}.$$

Example 21 Let G be a graph on t vertices with exactly two edges incident on four vertices (a matching with two edges), then

$$\mathcal{L}(G) = \frac{1}{t!} \sum_{i=2}^t \left(\frac{i-2}{i} + \frac{i-1}{i+1} + \dots + \frac{n-3}{n-1} \right).$$




FIG. 1: All non-isomorphic graphs on t vertices, where $t \leq 4$, and their likelihood.

By making use of Theorem 16, we can prove in a straightforward way that a graph and its complement have equal likelihood. The *complement* of a graph $G = (V, E)$, denoted by $\overline{G}$, is the graph such that $V(\overline{G}) = V(G)$ and $E(\overline{G}) = V(G) \times V(G) - \{\{v_i, v_i\} : v_i \in V(G)\} - E(G)$.

Proposition 22 Let G be any graph. Then $\mathcal{L}(G) = \mathcal{L}(\overline{G})$.

V. CONCLUSIONS

We have used a model of graph growth to introduce a notion of graph likelihood and we have then discussed some of its basic aspects. This is the probability that a graph is grown with the model. We have proposed an algorithm for the computation of the likelihood, and we have bounded this graph invariant in terms of the automorphism group. We conclude with two natural open problems:

Problem 23 How hard is to compute the likelihood?

Problem 24 Which graphs are extremal with respect to the likelihood?

Acknowledgments. We would like to thank Ginestra Bianconi, Sebi Cioaba, Chris Godsil, Anastasia Koroto, Matt DeVos, and Svante Janson.

-
- [1] C. Godsil, G. Royle, *Algebraic Graph Theory*, Springer-Verlag, 2004.
 - [2] S. Janson, S. Severini, An example of graph limits of growing sequences of random graphs, June 2012. [arXiv:1206.4586v1](https://arxiv.org/abs/1206.4586v1) [math.CO]
 - [3] Wikipedia contributors, “Infinite monkey theorem” *Wikipedia, The Free Encyclopedia*, http://en.wikipedia.org/wiki/Infinite_monkey_theorem (accessed February 21, 2013).